

DrugFacts

www.drugabuse.gov

Prescription Opioids

What are prescription opioids?

Opioids are a class of drugs naturally found in the opium poppy plant. Some prescription opioids are made from the plant directly, and others are made by scientists in labs using the same chemical structure. Opioids are often used as medicines because they contain chemicals that relax the body and can relieve pain. Prescription opioids are used mostly to treat moderate to severe pain, though some opioids can be used to treat coughing and diarrhea. Opioids can also make people feel very relaxed and “high” – which is why they are sometimes used for non-medical reasons. This can be dangerous because opioids can be highly addictive, and overdoses and death are common. Heroin is one of the world’s most dangerous opioids, and is never used as a medicine in the United States.



Photo by ©iStock.com/[Pictac](#)/

Popular slang terms for opioids include Oxy, Percs, and Vikes.

What are common prescription opioids?

- hydrocodone (Vicodin®) oxycodone (OxyContin®, Percocet®)
- oxymorphone (Opana®)
- morphine (Kadian®, Avinza®)
- codeine
- fentanyl

How do people misuse prescription opioids?

Prescription opioids used for pain relief are generally safe when taken for a short time and as prescribed by a doctor, but they can be misused. People misuse prescription opioids by:

- taking the medicine in a way or dose other than prescribed
- taking someone else’s prescription medicine
- taking the medicine for the effect it causes—to get high

When misusing a prescription opioid, a person can swallow the medicine in its normal form. Sometimes people crush pills or open capsules, dissolve the powder in water, and inject the liquid into a vein. Some also snort the powder.

How do prescription opioids affect the brain?

Opioids bind to and activate opioid receptors on cells located in many areas of the brain, spinal cord, and other organs in the body, especially those involved in feelings of pain and pleasure. When opioids attach to these receptors, they block pain signals sent from the brain to the body and release large amounts of dopamine throughout the body. This release can strongly reinforce the act of taking the drug, making the user want to repeat the experience.

What are some possible effects of prescription opioids on the brain and body?

In the short term, opioids can relieve pain and make people feel relaxed and happy. However, opioids can also have harmful effects, including:

- drowsiness
- confusion
- nausea
- constipation
- euphoria
- slowed breathing

Opioid misuse can cause slowed breathing, which can cause hypoxia, a condition that results when too little oxygen reaches the brain. Hypoxia can have short- and long-term psychological and neurological effects, including coma, permanent brain damage, or death.

Researchers are also investigating the long-term effects of opioid addiction on the brain, including whether damage can be reversed.

What are the other health effects of opioid medications?

Older adults are at higher risk of accidental misuse or abuse because they typically have multiple prescriptions and chronic diseases, increasing the risk of drug-drug and drug-disease interactions, as well as a slowed metabolism that affects the breakdown of drugs.

Sharing drug injection equipment and having impaired judgment from drug use can increase the risk of contracting infectious diseases such as HIV and from unprotected sex.

Prescription Opioids and Heroin

Prescription opioids and heroin are chemically similar and can produce a similar high. In some places, heroin is cheaper and easier to get than prescription opioids, so some people switch to using heroin instead. Nearly 80 percent of Americans using heroin (including those in treatment) reported misusing prescription opioids prior to using heroin.^{1,2}

However, while prescription opioid misuse is a risk factor for starting heroin use, only a small fraction of people who misuse pain relievers switch to heroin. This suggests that prescription opioid misuse is just one factor leading to heroin use. Read more about this intertwined problem in our [Prescription Opioids and Heroin Research Report](#).

Can I take prescription opioids if I'm pregnant?

If a woman uses prescription opioids when she's pregnant, the baby could develop dependence and have withdrawal symptoms after birth. This is called neonatal abstinence syndrome, which can be treated with medicines. Use during pregnancy can also lead to miscarriage and low birth weight. Read more in the [Substance Use in Women Research Report](#).

It can be difficult for a person with an opioid addiction to quit, but pregnant women who seek treatment have better outcomes than those who quit abruptly. Methadone and buprenorphine are the standard of care to treat opioid-dependent pregnant women. Methadone or buprenorphine maintenance combined with prenatal care and a comprehensive drug treatment program can improve many of the adverse outcomes associated with untreated opioid addiction. If a woman is unable to quit before becoming pregnant, treatment with methadone or buprenorphine during pregnancy improves the chances of having a healthier baby at birth.

In general, it is important to closely monitor women who are trying to quit drug use during pregnancy and to provide treatment as needed.

Can a person overdose on prescription opioids?

Yes, a person can overdose on prescription opioids. An opioid overdose occurs when a person uses enough of the drug to produce life-threatening symptoms or death. When people overdose on an opioid medication, their breathing often slows or stops. This can decrease the amount of oxygen that reaches the brain, which can result in coma, permanent brain damage, or death.

How can an opioid overdose be treated?

If you suspect someone has overdosed, the most important step to take is to call 911 so he or she can receive immediate medical attention. Once medical personnel arrive, they will administer naloxone. Naloxone is a medicine that can treat an opioid overdose when given right away. It works by rapidly binding to opioid receptors and blocking the effects of opioid drugs. Naloxone is available as an injectable (needle) solution, a hand-held auto-injector (EVZIO®), and a nasal spray (NARCAN® Nasal Spray).

Some states have passed laws that allow pharmacists to dispense naloxone without a personal prescription. This allows friends, family, and others in the community to use the auto-injector and nasal spray versions of naloxone to save someone who is overdosing.

Read more on our [Naloxone webpage](#).

Tolerance vs. Dependence vs. Addiction

Long-term use of prescription opioids, even as prescribed by a doctor, can cause some people to develop **a tolerance**, which means that they need higher and/or more frequent doses of the drug to get the desired effects.

Drug **dependence** occurs with repeated use, causing the neurons to adapt so they only function normally in the presence of the drug. The absence of the drug causes several physiological reactions, ranging from mild in the case of caffeine, to potentially life threatening, such as with heroin. Some chronic pain patients are dependent on opioids and require medical support to stop taking the drug.

Drug **addiction** is a chronic disease characterized by compulsive, or uncontrollable, drug seeking and use despite harmful consequences and long-lasting changes in the brain. The changes can result in harmful behaviors by those who misuse drugs, whether prescription or illicit drugs.

Can use of prescription opioids lead to addiction?

Yes, repeated misuse of prescription opioids can lead to a substance use disorder (SUD), a medical illness which ranges from mild to severe and from temporary to chronic. Addiction is the most severe form of an SUD. An SUD develops when continued misuse of the drug changes the brain and causes health problems and failure to meet responsibilities at work, school, or home.

People addicted to an opioid medication who stop using the drug can have severe withdrawal symptoms that begin as early as a few hours after the drug was last taken. These symptoms include:

- muscle and bone pain
- sleep problems
- diarrhea and vomiting
- cold flashes with goose bumps
- uncontrollable leg movements
- severe cravings

These symptoms can be extremely uncomfortable and are the reason many people find it so difficult to stop using opioids. There are medicines being developed to help with the withdrawal process, and the U.S. Food and Drug Administration (FDA) approved sale of a device, [NSS-2 Bridge](#), that can help ease withdrawal symptoms. The NSS-2 Bridge device is a small electrical nerve stimulator placed behind the person's ear, that can be used for up to five days during the acute withdrawal phase. There are also medicines being developed to help with the withdrawal process. The FDA approved lofexidine, a non-opioid medicine designed to reduce opioid withdrawal symptoms.

What type of treatment can people get for addiction to prescription opioids?

A range of treatments including medicines and behavioral therapies are effective in helping people with opioid addiction.

Two medicines, buprenorphine and methadone, work by binding to the same opioid receptors in the brain as the opioid medicines, reducing cravings and withdrawal symptoms. Another medicine, naltrexone, blocks opioid receptors and prevents opioid drugs from having an effect.

Behavioral therapies for addiction to prescription opioids help people modify their attitudes and behaviors related to drug use, increase healthy life skills, and persist with other forms of treatment, such as medication. Some examples include, cognitive behavioral therapy which helps modify the patient's drug use expectations and behaviors, and also effectively manage triggers and stress. Multidimensional family therapy, developed for adolescents with drug use problems, addresses a range of personal and family influences on one's drug use patterns and is designed to improve overall functioning. These behavioral treatment approaches have proven effective, especially when used along with medicines. Read more about drug addiction treatment in our [Treatment Approaches for Drug Addiction DrugFacts](#).

Points to Remember

- Prescription opioids are used mostly to treat moderate to severe pain, though some opioids can be used to treat coughing and diarrhea.
- People misuse prescription opioids by taking the medicine in a way other than prescribed, taking someone else's prescription, or taking the medicine to get high. When misusing a prescription opioid, a person may swallow, inject, or snort the drug.
- Opioids bind to and activate opioid receptors on cells located in the brain, spinal cord, and other organs in the body, especially those involved in feelings of pain and pleasure, and can strongly reinforce the act of taking the drug, making the user want to repeat the experience.
- People who use prescription opioids can feel relaxed and happy, but also experience drowsiness, confusion, nausea, constipation, and slowed breathing.
- Prescription opioids have effects similar to heroin. While prescription opioid misuse is a risk factor for starting heroin use, only a small fraction of people who misuse opioid pain relievers switch to heroin.
- A person can overdose on prescription opioids. Naloxone is a medicine that can treat an opioid overdose when given right away.
- Prescription opioid use, even when used as prescribed by a doctor can lead to a substance use disorder, which takes the form of addiction in severe cases. Withdrawal symptoms include muscle and bone pain, sleep problems, diarrhea and vomiting, and severe cravings.
- A range of treatments including medicines and behavioral therapies are effective in helping people with an opioid use disorder.

Learn More

For more information about opioid medications, see our:

- [Commonly Abused Drugs chart](#)
- [Misuse of Prescription Drugs Research Report](#)

This publication is available for your use and may be reproduced **in its entirety** without permission from the NIDA. Citation of the source is appreciated, using the following language:

Source: National Institute on Drug Abuse; National Institutes of Health; U.S. Department of Health and Human Services.

Updated June 2018

References

1. Muhuri PK, Gfroerer JC, Davies MC. *Associations of Nonmedical Pain Reliever Use and Initiation of Heroin Use in the United States*. Rockville, MD: Substance Abuse and Mental Health Services Administration; 2013. <http://archive.samhsa.gov/data/2k13/DataReview/DR006/nonmedical-pain-reliever-use-2013.pdf>. Accessed May 13, 2016.
2. Jones CM. Heroin use and heroin use risk behaviors among nonmedical users of prescription opioid pain relievers - United States, 2002-2004 and 2008-2010. *Drug Alcohol Depend*. 2013;132(1-2):95-100. doi:10.1016/j.drugalcdep.2013.01.007.